

Fax to 1-866-588-6940

PROLASTIN-C LIQUID Prescription and Enrollment Form/SMN

Patient Information

Patient Name (Last, First) _____
 Last four digits of Social Security # _____ Gender Male Female Date of Birth _____
 Address _____ Apt _____ City _____ State _____ Zip _____
 Home Phone _____ Best Time/Day _____ PROLASTIN DIRECT Hours of Operation: 7 am–7 pm CST
 Cell/Other Phone _____ Email Address _____

Please attach front and back copy of patient's insurance cards.

Medical Information

ICD-10 Diagnosis: Alpha₁-Antitrypsin Deficiency E88.01 Panlobular Emphysema J43.1 Other _____
 AAT Phenotype: _____ FEV₁ _____ % predicted DLCO _____ % predicted Serum AAT Level _____ mg/dL _____ μM
 Weight _____ lb *OR* _____ kg Allergies None *OR* Specify _____

Steps to e-Prescribe PROLASTIN-C LIQUID

1. Fax in the PROLASTIN-C LIQUID Prescription and Enrollment Form/SMN
2. Prescribe PROLASTIN-C LIQUID Intravenous Solution
3. Choose Dohmen Life Science Services (DLSS) as the dispensing pharmacy

Attach documentation of the following medical information:

History/Physical Summary Last Chest X-ray
 Most Recent PFTs Including FEV₁ AAT Phenotype or Genotype Lab Report

*DLSS is the exclusive dispensing pharmacy for PROLASTIN-C LIQUID

PROLASTIN-C LIQUID Prescription/Orders

PROLASTIN-C LIQUID (alpha₁-protease inhibitor [human]) Number of Refills: 12 months *Dispense up to a 30-day supply* Other Number of Refills _____
Dosage: 60 mg/kg (+/- 10%) IV weekly *OR* Other dose/frequency _____
Rate: As tolerated by patient up to 0.08 mL/kg/min *OR* Other rate _____

- Epinephrine Inj Auto 0.3 mg/0.3 mL. Inject from autodevice, prn, for allergic reaction
- EMLA or lidocaine 2.5%/prilocaine 2.5%: Apply 2.5 g over 20-25 cm² of skin surface at least 1 hour prior to puncture
- Premedication/Other orders: _____

Monitor patient including VS before, Q15 during, and 5 minutes post infusion
 Provide medical supplies, including syringes and needles, to safely administer prescribed medication
 Skilled nursing visits for medication administration, assessment, and teaching

Intravenous Access and Flush Orders:

- Peripheral IV line Normal saline: 3-5 mL before infusion and 3-5 mL after infusion *OR* Other _____
- Implanted Port/Central Line (each lumen) Normal saline: 5-10 mL before infusion and 5-10 mL after infusion
 Heparin: 100 units/mL, 5 mL after infusion, and _____ (frequency)

First Dose Location Preference(s): Home *OR* Medical facility (name, phone of preferred facility, if any): _____
 List of facilities where physician has privileges: _____

First Dose in Home Orders:

- Establish primary IV line with 250-500 mL of normal saline *OR* Other _____ at KVO rate prior to infusion
- Monitor patient including VS before, Q15 during, and 30 minutes post infusion
- Diphenhydramine 25-50 mg IV as directed for severe allergic reaction or anaphylaxis

Physician/Office Information

Physician Name (print) _____ Office Phone _____
 Address _____ Suite _____ Office Fax _____
 City _____ State _____ Zip _____
 Office Contact Person _____ Phone _____ NPI# _____
 Office Contact Person Email Address _____

By signing below, I certify that the therapy described above is medically necessary and that the information provided is accurate to the best of my knowledge. I also attest that I have obtained the patient's authorization to release the above information and such other personal information as may be necessary to PROLASTIN DIRECT, Dohmen Life Science Services, and/or their agents. If the patient is 18 years old or younger, I attest that I have obtained permission from the patient's legal guardian.

Physician Signature _____ Date _____
Substitution Permitted *Dispense as Written*

Fax completed form and documentation to 1-866-588-6940. To reach the PROLASTIN DIRECT team, call 1-800-305-7881.

Please see Important Safety Information on back and accompanying full Prescribing Information for PROLASTIN-C LIQUID.

Steps to e-Prescribe PROLASTIN-C LIQUID

1. Fax in the PROLASTIN-C LIQUID Prescription and Enrollment Form/SMN
 2. Prescribe PROLASTIN-C LIQUID Intravenous Solution
 3. Choose Dohmen Life Science Services (DLSS) as the dispensing pharmacy
- *DLSS is the exclusive dispensing pharmacy for PROLASTIN-C LIQUID

Documentation Checklist

Please fax the following documentation to the PROLASTIN DIRECT team so that we can start the insurance authorization process on your patient's behalf

1. Completed PROLASTIN-C LIQUID Prescription and Enrollment Form/SMN
2. Front and back copy of patient's insurance cards
3. History / Physical Summary
4. Last Chest X-ray diagnostic report
5. Most recent Pulmonary Function Report including FEV₁
6. AAT Lab Reports:
 - a) Serum AAT Level
 - b) AAT Phenotype or Genotype

* Note: If you have both genotype and phenotype results, you may provide both, but insurance plans usually only require one of the two.

Fax completed form and documentation to 1-866-588-6940. To reach the PROLASTIN DIRECT team, call 1-800-305-7881.

Important Safety Information

PROLASTIN®-C LIQUID is an alpha₁-proteinase inhibitor (human) (alpha₁-PI) indicated for chronic augmentation and maintenance therapy in adults with clinical evidence of emphysema due to severe hereditary deficiency of alpha₁-PI (alpha₁-antitrypsin deficiency).

Limitations of Use

- The effect of augmentation therapy with any alpha₁-PI, including PROLASTIN-C LIQUID, on pulmonary exacerbations and on the progression of emphysema in alpha₁-PI deficiency has not been conclusively demonstrated in randomized, controlled clinical trials
- Clinical data demonstrating the long-term effects of chronic augmentation or maintenance therapy with PROLASTIN-C LIQUID are not available
- PROLASTIN-C LIQUID is not indicated as therapy for lung disease in patients in whom severe alpha₁-PI deficiency has not been established

PROLASTIN-C LIQUID is contraindicated in immunoglobulin A (IgA)-deficient patients with antibodies against IgA or patients with a history of anaphylaxis or other severe systemic reaction to alpha₁-PI products.

Hypersensitivity reactions, including anaphylaxis, may occur. Monitor vital signs and observe the patient carefully throughout the infusion. If hypersensitivity symptoms occur, promptly stop PROLASTIN-C LIQUID infusion and begin appropriate therapy.

Because PROLASTIN-C LIQUID is made from human plasma, it may carry a risk of transmitting infectious agents, eg, viruses, the variant Creutzfeldt-Jakob disease (vCJD) agent, and, theoretically, the Creutzfeldt-Jakob disease (CJD) agent. This also applies to unknown or emerging viruses and other pathogens.

The most common adverse reactions during PROLASTIN-C LIQUID clinical trials in >5% of subjects were diarrhea and fatigue, each of which occurred in 2 subjects (6%).

Please see accompanying full Prescribing Information for PROLASTIN-C LIQUID.